

# ACCELERATING COSMOLOGIES WITH AN ANISOTROPIC EQUATION OF STATE

TOMI KOIVISTO

Helsinki Institute of Physics, P.O. Box 64, FIN-00014 Helsinki, Finland

DAVID F. MOTA

Institute for Theoretical Physics, University of Heidelberg, 69120 Heidelberg, Germany  
*Draft version February 1, 2008*

## ABSTRACT

If the dark energy equation of state is anisotropic, the expansion rate of the universe becomes direction-dependent at late times. We show that such models are not only cosmologically viable but that they could explain some of the observed anomalies in the CMB. The possible anisotropy can then be constrained by studying its effects on the luminosity distance-redshift relation inferred from several observations. A vector field action for dark energy is also presented as an example of such possibility.

*Subject headings:* (cosmology:) cosmic microwave background; cosmology: miscellaneous; cosmology: observations; cosmology: theory; cosmology: large-scale structure of universe

## 1. INTRODUCTION

It is not clear whether the large-angle anomalies in the observed cosmic microwave background are of a cosmological origin and not due to systematics (Eriksen et al. 2004; Land and Magueijo 2005; Copi et al. 2006). Nevertheless, it is tempting to associate the apparent statistical anisotropy with dark energy, since the anomalies occur at the largest scales, and these enter inside the horizon at the same epoch that the dark energy dominance begins.

The paramount characteristic of dark energy is its negative pressure. We examine then the possibility that this pressure varies with the direction. Then also the universal acceleration becomes anisotropic, and one would indeed see otherwise unexpected effects at the smallest multipoles of CMB. At the background level we will find a similar CMB pattern as in an universe which is ellipsoidal at the era of last scattering (Campanelli et al. 2006). We then show that the supernovae could be used to distinguish these different scenarios and to constrain the possible anisotropic properties of dark energy.

There are several motivations for anisotropic models of dark energy. For instance, many dark energy models are in principle compatible with the FLRW metric but exhibit anisotropic stresses at the perturbative level, including nonminimally coupled fields, viscous fluids and modified gravity models (Koivisto and Mota 2006; Schimd et al. 2006; Mota et al. 2007; Koivisto & Mota 2007a,b). It is possible that a more accurate description of such models should take into account the anisotropic effects on the background expansion, which then breaks the statistical isotropy of perturbations too (as appears to have happened in the CMB (Eriksen et al. 2004; Land and Magueijo 2005; Copi et al. 2006)). Also, considering dark energy as an effective description of a backreaction of sizable inhomogeneities in dark matter (Buchert 2007; Behrend, Brown & Robbers 2007), the validity of a perfect fluid description could seem dubious.

To have a general description of an anisotropic dark energy component, we consider a phenomenological parameterization of dark energy in terms of its equation of state ( $w$ ) and two skewness parameters ( $\delta$ ,  $\gamma$ )

and include also a coupling term ( $Q$ ) between dark energy and a perfect fluid (dark matter). Previous studies anisotropic dark energy have mainly considered the anisotropic properties of the inhomogeneous perturbations (Koivisto and Mota 2006; Battye and Moss 2006) whereas our approach here is to focus on a smooth cosmology with the anisotropic pressure field. Whereas in a previous approach the anisotropy was only weakly constrained (Mota et al. 2007), we find that in the present description only a narrow parameter range can survive the observational tests. Similar anisotropic inflation has been considered in the early universe (Burd and Lidsey 1991; Ford 1989). Therefore we focus on the present paper more on the dark energy era, where the qualitative differences are that matter cannot be neglected and that the cosmologies do not isotropize. As a proof of a concept, we write also an explicit field theory example where a vector field drives the anisotropic acceleration of the universe.

An anisotropic expansion is not compatible with the Robertson-Walker (RW) metric. Hence we use the B(I) (Bianchi type I) metric which generalizes the flat RW metric and may be employed to obtain limits on cosmological skew pressures from the CMB (Barrow 1997). The line-element of a B(I) universe is:

$$ds^2 = -dt^2 + a^2(t)dx^2 + b^2(t)dy^2 + c^2(t)dz^2. \quad (1)$$

There are thus three scale factors, and consequently three expansion rates. In principle all these could be different, and in the limiting case that all of them are equal one recovers the RW case. It is useful to express the mean expansion rate as an average Hubble rate  $H$

$$H \equiv \frac{1}{3} \left( \frac{\dot{a}}{a} + \frac{\dot{b}}{b} + \frac{\dot{c}}{c} \right), \quad (2)$$

(where an overdot means derivative wrt  $t$ ) and then the differences of the expansion rates as the Hubble-normalized shear  $R$  and  $S$  (Barrow 1997)

$$R \equiv \frac{1}{H} \left( \frac{\dot{a}}{a} - \frac{\dot{b}}{b} \right), \quad S \equiv \frac{1}{H} \left( \frac{\dot{a}}{a} - \frac{\dot{c}}{c} \right). \quad (3)$$

We consider a universe filled with a perfect fluid having the energy-momentum tensor  $T_{(m)\nu}^\mu = \text{diag}(-1, w_m, w_m, w_m)\rho_m$ , and dark energy, which we allow to have the most general energy-momentum tensor compatible with the metric (1)

$$T_{\nu}^{\mu} = \text{diag}(-1, w, w + 3\delta, w + 3\gamma)\rho. \quad (4)$$

The generalized Friedmann equation may be written as

$$H^2 = \frac{8\pi G}{3} \frac{\rho_m + \rho}{1 - \frac{1}{9}(R^2 + S^2 - RS)}. \quad (5)$$

We let the two components also interact. The continuity equations are then

$$\dot{\rho}_m + 3H(1 + w_m)\rho_m = QH\rho, \quad (6)$$

and

$$\dot{\rho} + 3 \left[ (1 + w)H + \delta \frac{\dot{b}}{b} + \gamma \frac{\dot{c}}{c} \right] \rho = -QH\rho, \quad (7)$$

where  $Q$  determines the coupling. If it vanishes, it follows that  $\rho_m \sim (abc)^{-1-w_m}$  and  $\rho \sim (abc)^{-1-w}b^{3\delta}c^{3\gamma}$ . Defining  $x \equiv \frac{1}{3} \log(-g)$ , where the metric determinant  $g = -abc$ , one notes that  $H = \dot{x}$ . We will use  $x$  as our time variable rather than  $t$ . Derivative wrt  $x$  is denoted by star. We also define the dimensionless density fractions

$$\Omega_m \equiv \frac{8\pi G}{3} \frac{\rho_m}{H^2}, \quad U \equiv \frac{\rho}{\rho_m + \rho}.$$

Using  $R$ ,  $S$ , and  $U$  as our dynamical variables, the system can finally be written as

$$\begin{aligned} U^* &= U(U - 1) [\gamma(3 + R - 2S) + \delta(3 - 2R + S) + 3(w - w_m)] - UQ, \\ S^* &= \frac{1}{6} (9 - R^2 + RS - S^2) \\ &\quad x \{S[U(\delta + \gamma + w - w_m) + w_m - 1] - 6\gamma U\}, \\ R^* &= \frac{1}{6} (9 - R^2 + RS - S^2) \\ &\quad x \{R[U(\delta + \gamma + w - w_m) + w_m - 1] - 6\delta U\}. \end{aligned} \quad (8)$$

Note the coupling term  $Q$  appears only in the evolution equation for  $U$ . Nevertheless its presence can change the dynamics completely (Koivisto and Mota 2008).

## 2. VECTOR FIELD

As a proof of concept, we present an explicit field theory for the anisotropically stressed dark energy. Consider the vector field action

$$S = \int d^4x \sqrt{-g} \left[ \frac{R}{16\pi G} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - V + qL_m \right] \quad (9)$$

where the kinetic term involves  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ . The potential  $V$  and the possible couplings  $q$  are understood as functions of the quadratic term  $A^2 = A_\mu A^\mu$ . The field equations are  $G_{\mu\nu} = 8\pi G (T_{\mu\nu}^m + T_{\mu\nu}^A)$ , where the energy-momentum tensor of the vector field follows by varying wrt the metric

$$T_{\mu\nu}^A = F_{\mu\alpha} F_{\nu}^{\alpha} + 2V' A_\mu A_\nu - \left( \frac{1}{4} F_{\alpha\beta} F^{\alpha\beta} + V \right) g_{\mu\nu}. \quad (10)$$

Prime denotes derivative wrt  $A^2$ . With the metric (1), the equation of motion for the  $A_0$  component of the field

dictates that  $[q'L_m + V']A_0 = 0$ . Thus we confine to purely spatial vector fields. The spatial components of the energy-momentum tensor are,

$$T_j^i A = \frac{1}{a_i^2} (-\dot{A}_i \dot{A}_j + 2V' A_i A_j) + \left( \frac{1}{2} \sum_{k=1}^3 \frac{A_k^2}{a_k^2} - V \right) \delta_j^i.$$

The off-diagonal terms should vanish. Thus one is restricted to consider only vectors which are parallel to one of the coordinate axis. There, however, could be many such fields.

In the RW case all the diagonal components of  $T_j^i$  should be equal, which indeed requires three vector fields, one in each coordinate direction, of exactly equal magnitudes, as the triads of Armendariz-Picon (Armendariz-Picon 2004). With the metric (1), the only constraint is that every vector field should be along a coordinate axis. Any system of these vector fields is thus *required* to have an anisotropic equation of state, unless it reduces to the very special triad case. Let us therefore briefly look at the case of a single vector field with  $A_1 = A_3 = 0$ . Notice that, in analogy to system (8), the coupling  $Q$  is now related to the function  $q$ . If we define the additional dimensionless parameters

$$X \equiv \frac{\dot{A}_2}{bH}, \quad Y \equiv \frac{V}{H^2}, \quad Y_1 = 2 \frac{V' A_2}{b^2 H^2},$$

one can relate the present case to the general notation by noting that  $H^2 \rho = X^2/2 + Y$ , and that if  $S = 0$ , then

$$w = \frac{X^2 - 2Y}{X^2 + 2Y}, \quad \delta = \frac{-X^2 - Y_1}{\frac{3}{2}X^2 + 3Y}. \quad (11)$$

We see that the anisotropy is naturally small if the field is either subdominant or near its minimum. Such models, even though perturbatively close to standard cosmology, would be excluded by imposing the RW symmetry. Viable models with large anisotropy do also exist as shown below. An extensive investigation of specific models will be undertaken elsewhere (Koivisto and Mota 2008).

## 3. SCALING SOLUTIONS (IN THE AXISYMMETRIC CASE)

To begin, we will assume for simplicity that 1) the perfect fluid is minimally coupled dark matter,  $w_m = Q = 0$ , and further that 2) the skewness together with the equation of state of dark energy is constant,  $\dot{w} = \dot{\delta} = \dot{\gamma} = 0$  (see (Koivisto and Mota 2008) for an extensive analysis with these assumptions relaxed). In our model the universe is then initially (close to) isotropic, but is driven to anisotropic expansion by the skewness of dark energy. Under the additional simplifying assumption of axial symmetry ( $S = \gamma = 0$ ), in FIG. 1 we indicate the final possible stages of such universe, corresponding to the fixed points of the system (8). The universe will then end up in three possible scenarios: (I) The isotropic RW case corresponding to

$$R = 0, \quad U = 0. \quad (12)$$

(II) An anisotropic dark energy dominated solution with

$$R = \frac{6\delta}{w + \delta - 1}, \quad U = 1. \quad (13)$$

Or (III) a scaling solution,

$$R = \frac{3(w + \delta)}{2\delta}, \quad U = \frac{-(w + \delta)}{3\delta^2 - 2\delta w - w^2}. \quad (14)$$




FIG. 1.— Asymptotic state from an Einstein-deSitter stage. The future fate depends on the dark energy properties  $w$  and  $\delta$  and is classified into three possibilities. If  $\delta + w > 0$ , the isotropically expanding dust domination (solution (12) for which  $U = 0$ ) continues forever. Otherwise, the universe will end up expanding anisotropically and either dominated by dark energy (solution (13) for which  $U = 1$ ) or exhibiting a scaling property (solution (14) for which  $0 < U < 1$ ).

Given that  $w$  is negative, this solution accelerates  $w_{eff} \equiv -\frac{2}{3}\frac{H^*}{H} - 1 < -\frac{1}{3}$ , if  $w < \delta < \frac{w}{3}$  or  $-\frac{w}{5} < \delta < -\frac{w}{3}$ . Note that in the  $R = 0$  case scaling solutions could only be found for coupled components.

Notice that within the RW universe it has been proven difficult to address the coincidence problem by finding a model entering from a matter-dominated scaling solution to an accelerating scaling solution. Allowing for the presence of three expansion rates opens up the possibility of describing a universe entering from a perfect fluid dominated scaling to an anisotropically accelerating scaling era. This might eventually help to understand the coincidence problem, since then matter and dark energy would have had similar energy densities both in the past and in the future (in the past the dark energy fraction, if constant should not exceed about 1/10 Doran and Robbers (2006)).

We now study the observational implications of models with nonzero skewness, relaxing the assumption 2). The relevant case is then a universe entering from RW solution (12) to an anisotropically accelerating universe (which generalizes (13) if  $\gamma \neq 0$ ).

#### 4. CMB ANISOTROPY

The B(I) background predicts a quadrupole pattern in the CMB. It is not clear how the fluctuations in the photon temperature distribution evolve when the background expansion becomes anisotropic, since the anisotropic sources, even if perturbatively small, couple to the perturbations at the first order (Koivisto and Mota 2008). It seems inevitable that statistically anisotropic features would then be, in principle, present in the whole spectrum of fluctuations (though in our case such features would be confined to larger scales as the anisotropy of the expansion be-

comes important only recently). Several large-angle anomalies have been reported in the CMB (Eriksen et al. 2004; Land and Magueijo 2005; Copi et al. 2006). It has been demonstrated that CMB spectrum of the tilted B(VIIh) background (Hawking 1968; Barrow et al. 1985; Pontzen and Challinor 2007) would have potential to model these anomalies (Jaffe et al. 2005) were it not in conflict with a dark energy dominated universe at the smaller angular scales (Jaffe et al. 2006).

The B(I) model reproduces the predictions of the concordance model at small scales, while featuring anomalies at large angles, since the (originally statistically isotropic) CMB field experiences an (statistically) anisotropic integrated Sachs-Wolfe effect in the nowadays ellipsoidal universe. Furthermore, it is plausible that these anomalies can match the observed ones, since a B(VIIh) model can be effectively described as a B(I) model with an additional anisotropic energy source<sup>1</sup>

As a first step, we check whether the background (1) is compatible with the available cosmological data. There are a number of calculations of CMB anisotropies in general Bianchi universes already available in the literature (Hawking 1968; Barrow et al. 1985; Pontzen and Challinor 2007). Here we consider the dark energy created quadrupole and the effects to the luminosity distance.

By considering the geodesic equation for photons, one can derive an equation for the redshift  $z$  of a photon arriving from the a direction  $\hat{\mathbf{p}}$

$$1 + z(\hat{\mathbf{p}}) = \frac{1}{a} \sqrt{1 + \hat{p}_y^2 e_y^2 + \hat{p}_z^2 e_z^2} \quad (15)$$

in terms of the eccentricities

$$e_y^2 = \left(\frac{a}{b}\right)^2 - 1, \quad e_z^2 = \left(\frac{a}{c}\right)^2 - 1. \quad (16)$$

Note that the scale factors and eccentricities here are evaluated at the time of last scattering in the case that the scale factors are all normalized to unity today. If  $T_*$  is the temperature at decoupling, the temperature field is given by  $T(\hat{\mathbf{p}}) = T_*/(1 + z(\hat{\mathbf{p}}))$ , and its spatial average is  $4\pi\bar{T} = \int d\Omega_{\hat{\mathbf{p}}} T(\hat{\mathbf{p}})$ . The anisotropy field is then

$$\frac{\delta T(\hat{\mathbf{p}})}{\bar{T}} = 1 - \frac{T(\hat{\mathbf{p}})}{\bar{T}}. \quad (17)$$

The coefficients in the spherical expansion of this anisotropy field are called  $a_{\ell m}$ , and due to orthogonality of spherical harmonics  $Y_{\ell m}$  are given by

$$a_{\ell m} = \int d\Omega_{\hat{\mathbf{p}}} \frac{\delta T(\hat{\mathbf{p}})}{\bar{T}} Y_{\ell m}^*. \quad (18)$$

The multipole spectrum is described by

$$Q_\ell = \sqrt{\frac{1}{2\pi} \frac{\ell(\ell+1)}{(2\ell+1)}} \sum_{m=-\ell}^{\ell} |a_{\ell m}|^2. \quad (19)$$

Expanding the redshifts (15) in powers of the eccentricities (16) one notes that the  $a_{\ell m}$  will be real (since there is only even dependence on the polar angle in the

<sup>1</sup> One may obtain the B(VIIIO) model from B(I) by adding anisotropic curvature. To obtain B(VIIh) model one should add also an isotropic curvature similar to such present in B(V) models.

anisotropy field and the imaginary parts of the  $e^{im\phi}$  integrate to zero), and that for all odd  $\ell$  the  $a_{\ell m}$  will vanish (since only even powers of the azimuthal angle appear in the expansion). To first order in  $e_{y,z}^2$ ,  $\bar{T} = aT_*[1 - \frac{1}{6}(e_z^2 + e_y^2)]$ , and in addition to the monopole, there is only the quadrupole

$$a_{20} = \frac{1}{3}\sqrt{\frac{\pi}{5}}(2e_z^2 - e_y^2), \quad a_{21} = a_{2-1} = 0, \quad (20)$$

$$a_{22} = a_{2-2} = -\sqrt{\frac{\pi}{30}}e_y^2 \Rightarrow Q_2 = \frac{2}{5\sqrt{3}}\sqrt{e_z^4 + e_y^4 - e_z^2e_y^2}$$

The observed value of  $Q_2$  is lower than the concordance model predicts (Hinshaw et al. 2007). It has been suggested in previous works that this discrepancy could be explained by an ellipsoidality of the universe (Campanelli et al. 2006; Jimenez and Maroto 2007). This would require that the anisotropy of the background is suitably oriented with respect to the intrinsic quadrupole and cancels its power to a sufficient amount. Too large anisotropy would of course only make the situation worse regardless of the orientation. Depending on the cosmological model, one should have  $Q_2 \lesssim 2 \cdot 10^{-5}$  to be consistent with observations taking into account the cosmic variance. The constraints this implies on the skewness of dark energy are very tight. However, we remark that in more general models, in particular with time-varying  $\delta$  and  $\gamma$ , one could allow more anisotropy. It is in principle possible for arbitrarily anisotropic expansion to escape detection from CMB (considering only the effects from the background), as long as the expansion rates evolve in such a way that  $e_z = e_y = 0$ . In other words, the (background) quadrupole vanishes, if each scale factor has expanded - no matter how anisotropically - the same amount since the last scattering. An example of such a scenario, derived from the action (9), is shown in FIG. 2.

## 5. SNIA LUMINOSITIES

The luminosity-redshift relationship of the SNIa data could be used to probe the possible anisotropies in the expansion history. This is a useful complementary probe since these objects are observed at the  $z < 2$  region, whereas CMB comes from much further away at  $z \sim 1100$ . The luminosity distance at the redshift  $z$  in the direction  $\hat{p}$  is now given by (Koivisto and Mota 2008)

$$d_L(z, \hat{p}) = (1+z) \int_{t_0}^{t(z)} \frac{dt}{\sqrt{\hat{p}_x^2 a^2 + \hat{p}_y^2 b^2 + \hat{p}_z^2 c^2}}. \quad (21)$$

To test this prediction with the data, we apply the formula (15) for each observed redshift of a supernova and match its luminosity distance inferred from the observation to the one computed from (21). In addition, we have to take into account also the angular coordinates of each individual supernovae in the sky to fix  $\hat{p}$  for each object. In our analysis we use the GOLD data set (Riess et al. 2006), which consists of five subsets of data<sup>2</sup>. We marginalize over the directions in the sky and over the present value of the Hubble constant. The results are

<sup>2</sup> SNIa Angular coordinates of each of the 182 GOLD supernovae can be found in (Riess et al. 2006; Astier et al. 2006) and <http://cfa-www.harvard.edu/ps/lists/Supernovae.html>




FIG. 2.— A minimally coupled vector field satisfying the quadrupole constraint. The solid (black) line is the dark energy density fraction  $U$ , the dashed (red) line is the effective equation of state of the universe  $w_{eff}$ , and the dash-dotted (blue) line describes the evolution of eccentricity,  $E = 500e_y^2$ . The potential is a double-power law  $V = mA^2 + \lambda A^{-4}$ . The dynamics of the field is such that though there are significant anisotropies, the eccentricity at the present is close to zero. This may be achieved with different power-law potentials, but requires fine-tuning of the mass scales  $m$  and  $\lambda$ .

summarized in FIGs. 3 and 4. The best-fit anisotropic models are only slightly preferred over the  $\Lambda$ CDM, the difference being  $\Delta\chi^2 \approx 1$ . Because of the additional parameters in the anisotropic models, the reasonable interpretation of these statistics is that the SNIa data favors isotropic expansion.




FIG. 3.— Limits on the (constant) skewness parameters of dark energy arising from the SNIa data, when  $\Omega_m = 0.3$  and  $w = -1$ . The contours correspond to 68.3, 90, 95.4 and 99 percent confidence limits.

The SNIa data constrains the skewness parameters much looser than the CMB quadrupole, if they are con-




FIG. 4.— Constraints arising from the SNIa in the axisymmetric case  $\gamma = 0$ . Inside the darker isosurface, the fit is as good as in the  $\Lambda$ CDM model,  $\chi^2 < 158$ . Inside the lighter isosurface, one has  $\Delta\chi^2 < 8.02$ . One notes that larger skewness  $\delta$  would typically be compatible with the SNIa data for phantom equations of state  $w < -1$  and large matter densities.

stant. However, in specific models with time-evolving  $\delta$  and  $\gamma$ , the constraints from CMB and from SNIa can be of comparable magnitude and allow anisotropy to an interesting degree. This means that the even if the CMB formed isotropically at early time, it could be distorted by the acceleration of the later universe in such

<sup>3</sup> Generating these effects at low redshift has an advantage that it relaxes constraints which would otherwise come from the CMB polarization (Pontzen and Challinor 2007) and could be strong for a given temperature anisotropy in isotropizing models because of

a way that it appears to us anomalous at the largest scales<sup>3</sup>. The future SNIa data, with considerably improved error bars (e.g. from the SNAP experiment, <http://snap.lbl.gov/>) on  $d_L(z, \hat{\mathbf{p}})$  might be used to rule out this possibility, and to distinguish whether the possible statistical anisotropy was already there at last scattering or whether it is due to dark energy.

## 6. CONCLUSIONS

In conclusion, dark energy with an anisotropic equation of state might be the culprit for both the cosmic acceleration and the large-angle anomalies in the CMB. This might also be the key to understand the coincidence problem. The present SNIa data allows anisotropic acceleration, but SNAP could set things straight about the skewness of dark energy and so of its nature. Such possibility would open a completely new window not only on the nature of the CMB anomalies but also into high energy physics models beyond the usual isotropic candidates of dark energy such as scalar fields or the cosmological constant.

We thank Y. Gaspar, H. Sigbjorn and the anonymous referee for useful comments. TK acknowledges support from the Magnus Ehrnrooth Foundation, the Finnish Cultural Foundation and EU FP6 Marie Curie Research and Training Network "UniverseNet" (MRTN-CT-2006-035863). DFM acknowledges support from the A. Humboldt Foundation and the Research Council of Norway through project number 159637/V30.

the significant polarization anisotropy at last scattering. However, anisotropic dark energy could evade this since the optical depth to  $z \sim 1$  is very small.

## REFERENCES

- Astier P. et al., *Astron. Astrophys.* **447**, 31 (2006).  
 Armendariz-Picon C., *JCAP* **0407**, 007 (2004).  
 Barrow J.D., *Phys. Rev.* **D55**, 7451 (1997).  
 Barrow et al., *Mon. Not. Roy. Astr. Soc.* **213**, 917 (1985).  
 R. A. Battye and A. Moss, *Phys. Rev. D* **74** (2006) 041301 [arXiv:astro-ph/0602377].  
 Behrend J., Brown I. and Robbers G., arXiv:0710.4964 [astro-ph].  
 Buchert T., *AIP Conf. Proc.* **910**, 361 (2007).  
 A. B. Burd and J. E. Lidsey, *Nucl. Phys. B* **351**, 679 (1991).  
 Campanelli L. et al., *Phys. Rev. Lett.* **97**, 131302 (2006).  
 Copi C. et al., *Mon. Not. Roy. Astr. Soc.* **367**, 79 (2006).  
 Doran M. and Robbers G., *JCAP* **0606**, 026 (2006).  
 Eriksen H.K. et al., *Astrophys. J.* **605**, 14 (2004).  
 L. H. Ford, *Phys. Rev. D* **40** (1989) 967.  
 S. W. Hawking, *Mon. Not. Roy. Astron. Soc.* **142** (1969) 129.  
 Hinshaw G. et al., *Astrophys. J. Suppl.* **170**, 288 (2007).  
 Jaffe T.R. et al., *Astrophys. J.* **629**, L1 (2005).  
 Jaffe T.R. et al., *Astrophys. J.* **644**, 701 (2006).  
 Jimenez and Maroto, *Phys. Rev.* **D76**, 023003 (2007).  
 T. Koivisto and D. F. Mota, *Phys. Rev. D* **73** (2006) 083502 [arXiv:astro-ph/0512135].  
 Koivisto T. and Mota D.F., (2007a) *Phys. Lett. B* **644**, 104 ;  
 Koivisto T. and Mota D.F., (2007b) *Phys. Rev. D* **75**, 023518;  
 T. Koivisto and D. F. Mota (2008), Work in progress.  
 Land and Magueijo, *Phys. Rev. Lett.* **95**, 071301 (2005).  
 Mota D. F., Kristiansen J., Koivisto T. and Groeneboom N., arXiv:0708.0830 [astro-ph].  
 A. Pontzen and A. Challinor, arXiv:0706.2075 [astro-ph].  
 Riess A.G. et al. (2006), astro-ph/0611572.  
 Schimd et al., *Phys. Rev. D* **71**, 083512 (2005).